

1-3-2000

Comment on “Inelastic Collapse of a Randomly Forced Particle” by Cornell et al.

J. Florencio

F. C. Sá Barreto

O. F. de Alcantara Bonfim
University of Portland, bonfim@up.edu

Follow this and additional works at: http://pilotscholars.up.edu/phy_facpubs

 Part of the [Physics Commons](#)

Citation: Pilot Scholars Version (Modified MLA Style)

Florencio, J.; Sá Barreto, F. C.; and Bonfim, O. F. de Alcantara, "Comment on “Inelastic Collapse of a Randomly Forced Particle” by Cornell et al." (2000). *Physics Faculty Publications and Presentations*. 45.
http://pilotscholars.up.edu/phy_facpubs/45

This Other is brought to you for free and open access by the Physics at Pilot Scholars. It has been accepted for inclusion in Physics Faculty Publications and Presentations by an authorized administrator of Pilot Scholars. For more information, please contact library@up.edu.

Comment on “Inelastic Collapse of a Randomly Forced Particle”

In their Letter [1] Cornell *et al.* investigated the dynamics of a particle in a one-dimensional box of length l subject to a random force $\eta(t)$. The particle motion is governed by the equation

$$\frac{d^2x}{dt^2} = \eta(t), \quad (1)$$

where $\eta(t)$ is a Gaussian noise with zero mean and a delta-correlated time dependence $\langle \eta(t)\eta(t') \rangle = 2\delta(t-t')$. The collisions with the boundaries are inelastic, with coefficient of restitution r ; that is, whenever the particle collides with a wall it rebounds with a speed which is r times smaller than the speed before collision. The authors report a remarkable transition in which below a critical value, $r_c = \exp(-\pi/\sqrt{3})$, the particle adheres to one of the walls after colliding with it an infinite number of times in a finite time. For $r > r_c$ the motion is ergodic. They also find that the value of r_c is independent of the box size and of the magnitude of a viscous damping term added to the equation of motion, Eq. (1). Through a series of rescaling of the position and time variables and followed by a transformation onto a stationary Gaussian process, the equation of motion is mapped onto that of a Brownian particle subject to a harmonic force, Eq. (6) of [1]. The critical value r_c was obtained from a mean-field analysis. They also performed a numerical integration of Eq. (6) and, upon transforming back onto the particle's original coordinates, they found evidence for the inelastic collapse for the particle trajectories, as predicted by their mean-field calculation. Numerical results for the collapse transition were displayed in Fig. 2 of their Letter.

We decided to numerically integrate Eq. (1) directly, in the hope to reproduce the results of Cornell *et al.* To our surprise we did not find *any instance of a collapse transition* for any values of r . In order to assert the robustness of the solutions, we used both second- and third-order stochastic Runge-Kutta methods [2] to integrate Eq. (1) for a variety of initial conditions on the positions and velocities. We used integration time steps in the interval 10^{-2} to 10^{-4} . We tested our numerical integration method against the exact result for both the mean square displacement and the mean square velocity of a Brownian particle. The agreement was within 2% or less of the exact values for 10^4 samples and times up to 150. In Fig. 1 we show our results for the trajectories of a particle with position $0 \leq x \leq 1$ and times up to $t = 100$ (same range of parameters used in Fig. 2 of [1]), for two values of the coefficient of restitution below and above the expected critical value $r_c = 0.163$. Clearly, no collapse transition is observed. We also find that the presence of a viscous damping term in Eq. (1)




FIG. 1. Trajectories of a particle confined in a one-dimensional box subject to a random force. The trajectories correspond to values of the coefficient of restitution above and below $r_c = 0.163$. No evidence of the collapsed phase is observed. Numerical integrations were performed using a third-order stochastic Runge-Kutta method with integration time step 10^{-2} , and initial conditions $(x, v) = (0, 1)$.

does not induce a collapse transition for *any* values of r and coefficient of friction.

We further analyze the model by calculating the power spectrum of $x(t)$. Our analysis shows that the averaged power spectrum is independent of the coefficient of restitution r . In addition, we find that the system exhibits two distinct $1/f^\alpha$ behaviors. For high frequencies we obtain a $1/f^2$ behavior for the power spectrum, as one would expect, which reflects the Brownian character of the particle motion at short times. On the other hand, at low frequencies we find a $1/f^4$ behavior for the power spectrum which can be attributed to the multiple collisions of the particle with the walls. We also find that in the low frequency regime the power α decreases as the damping coefficient increases. Details of these results will be presented elsewhere.

This work was partially supported by CNPq, FINEP, FAPEMIG, and MCT (Brazilian agencies).

J. Florencio and F.C. Sá Barreto

Departament de Física, UFMG
CP 702, 30123-970 Belo Horizonte, Brazil

O.F. de Alcantara Bonfim

Department of Physics, Reed College
Portland, Oregon 97202

Received 13 July 1999

PACS numbers: 05.40.Jc

- [1] S. J. Cornell, M. R. Swift, and A. J. Bray, Phys. Rev. Lett. **81**, 1142 (1998).
- [2] E. Helfand, Bell Syst. Tech. J. **56**, 2289 (1979); H. S. Greenside and E. Helfand, Bell Syst. Tech. J. **60**, 1927 (1981).